

Q-Wave Infarction Versus S-T Infarction

Nonspecificity of Electrocardiographic Criteria for Differentiating Transmural and Nontransmural Lesions

DAVID H. SPODICK, MD, DSc

Nonspecificity of the Electrocardiogram

For several decades cardiologists have considered abnormal Q waves synonymous with transmural myocardial infarction, as distinguished from infarcts characterized only by ST-T abnormalities, labelled "nontransmural" or "subendocardial." Curiously, journal reviewers have not vetoed these misnomers despite repeated marshalling of evidence demonstrating at autopsy that abnormal Q waves occur frequently with anatomically nontransmural infarction and may be absent with transmural infarcts.¹⁻³ This evidence first appeared about 40 years ago and has consistently been substantiated.⁴⁻¹⁹ Indeed, early investigators who reported an electrocardiographic distinction between the 2 anatomic types of infarct failed to prove it on examination of their own illustrations: among 4 electrocardiograms illustrating nontransmural infarcts, 1 shows incomplete left bundle branch block and 3 clearly have pathologic Q waves¹⁵; among 6 subendocardial infarcts demonstrated pathologically, 2 electrocardiograms show incomplete left bundle branch block, 2 have abnormal (≥ 0.04 second) Q waves in lead aVF, and 2 were too poorly reproduced to decipher.⁵ (This is especially surprising considering that illustrations invariably put the author's "best foot forward.") More recently, among 12 of 24 autopsied subjects with nontransmural infarcts, 10 had QRS changes, 8 with abnormal Q waves (including anatomically small infarcts); the other 12 patients had transmural infarcts, but 5 had no abnormal Q waves.⁸ Another recent report confirms postmortem documentation of transmural infarcts with only ST-T changes.¹¹ In a series of 154 autopsy subjects, most of those with nontransmural infarction had typical QRS changes and many with transmural infarcts lacked pathologic Q waves.⁷

It is possible that some earlier work may have established an *idée fixe* in the mind of cardiologists, despite the early availability of a mass of contrary evi-

dence.^{4-6,17,19} Indeed, Wilson, a pioneer in cardiology, and his colleagues¹⁹ demonstrated abnormal Q waves in nontransmural infarction both experimentally and clinically. The perceptive observations of Pipberger and Lopez²⁰ suggest that the erroneous criteria may have been reinforced by Prinzmetal's experiments of 1954, subsequently invalidated by the same investigator in 1957.

Conceptual Errors Inhibit Investigation

Whatever the reason for continued confusion regarding their anatomic extent, the pseudospecificity of Q and S-T myocardial infarctions remains a lasting myth. Indeed, investigators of the highest intellect and skill—despite explicitly acknowledging the material cited in References 1 to 3,²¹—write "regardless of the terms used. . .,"²¹ implying that slipshod terminology is unimportant. Such disregard by distinguished colleagues is disturbing. We are not dealing with minor matters such as "ECG" (English) versus "EKG" (Dutch and German). Arguments over terminology would be mindless quibbles if misleading labels did not perpetuate misleading concepts. False terminology does more than semantic damage: it does conceptual damage. If we continue to accept a Q infarct as dependably indicating a transmural lesion, we can inhibit consideration of what it really does mean. Indeed, do Q infarcts and S-T infarcts have distinctive gross anatomy? Given the ultimate nonspecificity of many electrocardiographic findings it should not be surprising that they overlap to such a degree that "transmural" and "nontransmural" are gross misnomers.

Q Infarcts Versus S-T Infarcts: Differences and Similarities

Although their gross anatomy cannot yet be predicted electrocardiographically, Q-wave and S-T infarcts differ clinically and physiologically. Prodromal symptoms are more frequent with S-T infarcts²²; vomiting is more frequent with Q-wave infarcts.¹⁸ Q-wave infarction implies more damage as measured by peak creatine phosphokinase levels,²³ with a greater tendency to undergo expansion.³⁴ Positron emission tomography²⁵ shows defects with homogeneous depression of tracer

From the Department of Medicine, University of Massachusetts Medical School, and the Division of Cardiology, St. Vincent Hospital, Worcester, Massachusetts. Manuscript received October 12, 1982; revised manuscript received October 28, 1982, accepted November 22, 1982.

Address for reprints: David H. Spodick MD, Director of Cardiology, St. Vincent Hospital, 25 Winthrop Street, Worcester, Massachusetts 01604.

TABLE I Q-Wave Infarction Versus S-T Infarction: Differences

	Q-Wave Infarct	S-T Infarct*
Fresh thrombosis	More	Less
Collateral vessels	Fewer	More
Damage		
↑ Enzyme levels	Higher	Lower
↓ Tracer concentration (PET)	Homogeneous	Unequal
LV wall abnormality,†	More	Less
improved by bypass	Less or none	More
Tendency to expand	More	Less
Prodromal symptoms	Less	More
Vomiting	More	Less
Atrioventricular and intraventricular blocks	More	Fewer
Congestive failure	More	Less
Early mortality	More	Less
Recurrent infarction	Less	More

* Without QRS abnormality. † Some studies differ.
PET = positron emission tomography.

accumulation in all Q-wave infarcts as compared with unequal tracer distribution in 23 of 24 S-T infarcts, with tomographic infarct size greater and residual tracer more depressed in Q-wave infarcts.²⁸ Two-dimensional echocardiograms showed more (100%) regional wall abnormality during Q-wave than S-T infarction (86%).^{26,27} Although total mortality may equalize over 5 to 10 years in the 2 groups, early mortality both in the hospital and after discharge is greater with Q-wave infarcts, whereas recurrent infarction and late mortality are much greater with S-T infarcts.^{28,30}

Although the number of diseased coronary arteries and the degree of occlusion throughout the system appear equal in both types,²³ fresh coronary thrombosis may be slightly³¹ to moderately¹⁰ more frequent with Q-wave infarction. Collateral vessels were demonstrated in 93% of all patients with S-T infarct and 78% with Q-wave infarct who had postinfarction angina, but in only 35% of patients with Q-wave infarct and no subsequent angina.³¹ Coronary bypass tends to improve wall motion in S-T infarcts but only in Q-wave infarcts in which transmural fibrosis is absent.^{32,33} Some investigators found reinfarction, angina, and late ventricular arrhythmias to occur equally in Q-wave and S-T infarcts, but more akinesis in patients with anterior Q-wave than with anterior S-T infarcts.³⁴ Yet, other investigators report comparable ejection fractions and wall motion abnormalities.²³ Finally, in a carefully matched, although retrospective, series S-T infarcts produced less congestive failure, fewer intraventricular conduction defects, and fewer atrial tachyarrhythmias; hypotension and ventricular arrhythmias occurred equally.²¹ Yet, the same study also reported a statistically equal occurrence of pericarditis.²¹ Because meticulous postmortem examination shows that in fresh infarction acute pericarditis always indicates transmural lesions,¹⁰ those non-Q-wave infarcts with pericarditis, although labelled "nontransmural,"²¹ certainly represented transmural infarction—more evidence of the electrocardiogram's anatomic non-specificity.

TABLE II Q-Wave Infarction Versus S-T Infarction: Similarities

Both either transmural or nontransmural
Number of diseased vessels
Degree of occlusion throughout coronary arteries
Ejection fraction*
Hypotension
Ventricular arrhythmia*
Late (>2 year) mortality

* Some studies differ.

Subdivisions Within Q Infarctions and S-T Infarctions

Investigations pertinent to this analysis correlated anatomically transmural and nontransmural infarcts with electrocardiographically Q-wave and S-T infarcts, irrespective of regional distribution.⁴⁻¹⁹ Fresh S-T infarcts usually occur with a normal QRS, but may also occur in patients with previous infarction, ventricular hypertrophy, preexcitation, or conduction disturbances producing QRS changes that mask or cancel processes producing new Q waves. Although most Q-wave infarcts are predominantly either inferior or anterior, it is not established whether either location is more likely to be transmural or nontransmural. Yet, location may have clinical importance: overall mortality in some series is worse with anterior infarcts, perhaps because of their tendency to be larger than inferior infarcts.^{35,36} In contrast, in the AMIS Study (4,524 participants) total mortality did not differ among infarct locations, including patients who lost abnormal Q waves.³⁷ Yet, ventricular dysfunction tends to be worse with anterior Q waves.^{38,39} Finally, both bundle branch block and significant myocardial failure are more frequent with anterior Q-wave infarcts, whereas sinus bradycardia, atrioventricular block, and atrial arrhythmias are more frequent with inferior Q-wave infarcts.²¹

Q-Wave Infarcts Versus S-T Infarcts

Tables I and II list the similarities and the many differences between Q-wave and S-T infarction. Q-wave infarcts include posterior infarction appearing as a new anterior R wave.

Conclusions: The unfortunate practice of mislabelling Q-wave infarction as "transmural" and S-T infarction as "nontransmural" does not correspond to the pathologic evidence. Q-wave infarcts and S-T infarcts differ clinically, physiologically, and prognostically but cannot be anatomically differentiated by electrocardiography. As with a variety of other loose or erroneous terminologies, the absence of precise labelling interferes with clarity of concept and may inhibit further investigation.⁴⁰ When a myocardial infarct is characterized strictly by electrocardiography, it should be described by the only accurate terminology—Q-wave infarct or S-T infarct. Confucius tells us:

If language is not used rightly, then what is said is not what is meant. If what is said is not what is meant, then that which ought to be done is left undone.

References

1. Spodick DH. Transmural vs. nontransmural infarction (letter). *Circulation* 1980;62:447-448.
2. Spodick DH. Anatomic nonspecificity of Q-wave and ST infarcts (letter). *Chest* 1981;79:245.
3. Spodick DH. Left ventricular thrombosis after myocardial infarction (letter). *N Engl J Med* 1982;305:1416.
4. Jaffe HL. Typical and atypical electrocardiograms in myocardial infarction caused by acute coronary occlusion and coronary insufficiency. *Dis Chest* 1955;27:243-250.
5. Levine HD, Ford RV. Subendocardial infarction: report of six cases and critical survey of the literature. *Circulation* 1950;1:246-262.
6. Master AM, Gubner R, Dack S, Jaffe HL. Differentiation of acute coronary insufficiency with myocardial infarction from coronary occlusion. *Arch Intern Med* 1941;67:647-655.
7. Raunio H, Rissanen V, Romppanen T, Jokinen Y, Rehnberg S, Heikin M, Pyörälä K. Changes in the QRS complex and ST segment in transmural and subendocardial infarctions. A clinicopathologic study. *Am Heart J* 1979;98:176-183.
8. Savage RM, Wagner GS, Ideker RE, Podolsky SA, Hackett DB. Correlation of postmortem anatomic findings with electrocardiographic changes in patients with myocardial infarction. *Circulation* 1977;55:279-285.
9. Madias JE, Chahine RA, Gorlin R, Blacklow DJ. A comparison of transmural and nontransmural acute myocardial infarction. *Circulation* 1974;49:498-504.
10. Erhardt LR. Clinical and pathological observations in different types of acute myocardial infarction. A study of 84 patients deceased after treatment in coronary care unit. *Acta Med Scand* 1974;Suppl 560:1-78.
11. Abbott JA, Scheinman MM. Nondiagnostic electrocardiogram in patients with acute myocardial infarction. *Am J Med* 1973;55:608-611.
12. Horan LG, Flowers NC, Johnson JC. Significance of the diagnostic Q wave of myocardial infarction. *Circulation* 1971;43:428-432.
13. Wilkinson RS, Schaefer A, Abildskov JA. Electrocardiographic and pathologic features of myocardial infarction in man. *Am J Cardiol* 1963;11:24-30.
14. Georas CS, Dahlquist E, Cutts FB. Subendocardial infarction. Correlation of clinical, electrocardiographic and pathologic data in 17 cases. *Arch Intern Med* 1963;111:488-492.
15. Cook RW, Edwards JE, Pruitt RD. Electrocardiographic changes in acute subendocardial infarction. I. Large subendocardial and large nontransmural infarcts. *Circulation* 1958;18:603-610.
16. Cook RW, Edwards JE, Pruitt RD. Electrocardiographic changes in acute subendocardial infarction. II. Small subendocardial infarcts. *Circulation* 1958;18:613-618.
17. Myers GB, Sears CH, Hiratzka T. Correlation of electrocardiographic and pathologic findings in ring like subendocardial infarction of left ventricle. *Am J Med Sci* 1951;222:417-421.
18. Ingram DA, Fulton RA, Portal RW, Aber CP. Vomiting as a diagnostic aid in acute ischemic cardiac pain. *Br Med J* 1980;281:636-637.
19. Wilson FN, Jonston FD, Rosenbaum FF, Erlanger H, Kossmann CE, Hecht H, Cotrim N, Menezes de Oliveira R, Scarsi R, Barker PS. The precordial electrocardiogram. *Am Heart J* 1944;27:19-31.
20. Pipberger HV, Lopez EA. "Silent" subendocardial infarcts. *Am Heart J* 1980;100:597-599.
21. Hutter AM, DeSanctis RW, Flynn T, Yeatman LA. Nontransmural myocardial infarction: a comparison of hospital and late clinical course of patients with that of matched patients with transmural anterior and transmural inferior myocardial infarction. *Am J Cardiol* 1981;48:595-602.
22. Scheidt S, Wolk M, Killip T. Unstable angina pectoris. Natural history, hemodynamics, uncertainties of treatment and the ethics of clinical study. *Am J Med* 1976;60:409-417.
23. Schulze RA, Pitt B, Griffith LSC, Ducci HH, Achuff SC, Baird MG, Humphries J. Coronary angiography and left ventriculography in survivors of transmural and nontransmural myocardial infarction. *Am J Med* 1978;64:108-113.
24. Hutchins GM, Bulkley BH. Infarct expansion versus extension: two different complications of acute myocardial infarction. *Am J Cardiol* 1978;41:1127-1132.
25. Geltman EM, Biello D, Welch MJ, Ter-Pogossian MM, Roberts R, Sobel BE. Characterization of nontransmural myocardial infarction by positron-emission tomography. *Circulation* 1982;65:747-755.
26. Horowitz RS, Morganroth J. Immediate detection of early high-risk patients with acute myocardial infarction using two-dimensional echocardiography evaluation of left ventricular regional wall motion abnormalities. *Am Heart J* 1982;103:814-818.
27. Horowitz RS, Morganroth J, Parrotto C, Chen CC, Soffer J, Pauletto FJ. Immediate diagnosis of acute myocardial infarction by two-dimensional echocardiography. *Circulation* 1982;65:323-329.
28. Marmor A, Sobel BE, Roberts R. Factors presaging early recurrent myocardial infarction ("extension"). *Am J Cardiol* 1981;48:603-610.
29. Cannon DS, Levy W, Cohen LS. The short- and long-term prognosis of patients with transmural and nontransmural myocardial infarction. *Am J Med* 1976;61:452-458.
30. Geltman EM, Ehsani AA, Campbell MK, Schiechtman K, Roberts R, Sobel BE. The influence of location and extent of myocardial infarction on long-term ventricular dysrhythmia and mortality. *Circulation* 1979;60:805-814.
31. Fuster V, Frye RL, Kennedy MA, Connolly DC, Mankin HT. The role of collateral circulation in the various coronary syndromes. *Circulation* 1979;59:1137-1142.
32. Roberts AJ, Moran JM, Sanders JH, Michaelis LL, Spies SM. Timing of myocardial revascularization following acute myocardial infarction (abstr). *Circulation* 1980;62:Suppl III:III-94.
33. Franklin K, Sutton M St. J, Mermelstein LJ, Cohen M, Harken A, Reichet N. Effect of left anterior descending coronary bypass on anterior wall function (abstr). *Circulation* 1982;II:II-90.
34. Taylor GJ, Humphries J, Mellits ED, Pitt B, Schulze RA, Griffith LSC, Achuff SC. Predictors of clinical course, coronary anatomy and left ventricular function after recovery from acute myocardial infarction. *Circulation* 1980;62:960-970.
35. Goldberg RJ, Kennedy HL, Szklo M, Tonascia J. Prognosis of anterior myocardial infarction revisited (abstr). *Am J Cardiol* 1979;43:370.
36. Geltman EM, Ehsani AA, Campbell MK, Roberts R, Sobel BE. Determinants of prognosis after initial subendocardial compared to transmural myocardial infarction: the importance of infarct size (abstr). *Am J Cardiol* 1979;43:370.
37. Wasserman AG, Bren GB, Ross AM, Richardson DW, Hutchinson RG, Rios JC. Prognostic implications of diagnostic Q waves after myocardial infarction. *Circulation* 1982;65:1451-1455.
38. Bodenheimer MM, Banka VS, Helfant RH. Q waves and ventricular asynergy: predictive value and hemodynamic significance of anatomic localization. *Am J Cardiol* 1975;35:615-618.
39. Palmeri ST, Harrison DG, Upton MT, Morris KG, Wagner GS. The electrocardiographic prediction of left ventricular ejection fraction following myocardial infarction (abstr). *Circulation* 1979;60:II-187.
40. Spodick DH. Cardioloecution. *Am J Cardiol* 1981;48:973-974.